# ReadyStream<sup>®</sup> Media Bag GranuCult<sup>®</sup> Buffered Peptone Water

Acc. ISO 6579, ISO 19250, ISO 21528, ISO 22964, ISO 6887, FDA-BAM and EP, irradiated

#### **Intended Use**

For the preliminary non-selective enrichment of bacteria, particularly pathogenic Enterobacteriaceae such as *Salmonella* and *Cronobacter*, from food and animal feed, water and other materials.

Each ReadyStream<sup>®</sup> media bag contains granulated Buffered Peptone Water broth for the generation of a liquid 10-fold stock concentrate. The ReadyStream<sup>®</sup> media bags are only to be used on the ReadyStream<sup>®</sup> instrument in combination with the ReadyStream<sup>®</sup> filter set for broth stock reconstitution and final dispensing of a diluted portion of the media.

#### **Mode of Action**

The broth is rich in nutrients and produces high resuscitation rates for sublethal injured bacteria and intense growth. The phosphate buffer system prevents bacterial damage caused by changes in the pH of the medium. Peptone including enzymatic digest of casein acts as a source of carbon, nitrogen, vitamins, and minerals whilst sodium chloride maintains the osmotic balance.

For *in vitro* use only.

#### **Typical composition**

Specified by ISO 6579, ISO/FDIS 6579-1, ISO 19250, ISO 21528, ISO 22964		Specified by FDA-BAM M192		Specified by EP 2.6.31, ISO 6785   IDF 93		ReadyStream <sup>®</sup> Media Bag GranuCult <sup>®</sup> Buffered Peptone Water acc. ISO 6579, ISO 21528, ISO 22964, FDA-BAM and EP, irradiated	
Enzymatic Digest of Casein*	10 g/L	Peptone	10 g/L	Peptone	10 g/L	Peptone (includes Enzymatic Digest of Casein)	10 g/L
NaCl	5 g/L	NaCl	5 g/L	NaCl	5 g/L	NaCl	5 g/L
$Na_2HPO_4$ x 12 H <sub>2</sub> O	9 g/L	$Na_2HPO_4^{**}$	3.5 g/L	$Na_2HPO_4$ x 12 H <sub>2</sub> O	9 g/L	$Na_2HPO_4$ x 12 H <sub>2</sub> O	9 g/L
KH <sub>2</sub> PO <sub>4</sub>	1.5 g/L	$KH_2PO_4$	1.5 g/L	KH <sub>2</sub> PO <sub>4</sub>	1.5 g/L	KH <sub>2</sub> PO <sub>4</sub>	1.5 g/L
Water	1000 mL/L	Water	1000 mL/L	Water	1000 mL/L	Water	n/a
pH at 25 °C	7.0 ± 0.2	pH at 25 °C	7.0 ± 0.2	pH at 25 °C	7.0 ± 0.2	pH at 25 °C	7.0 ± 0.2

\* IS0/FDIS 6579-1 specifies: Peptone - for example, enzymatic digest of casein. \*\*  $3.57 \text{ g} \text{ Na}_{2}\text{HPO}_{4}$  anhydrous is equivalent to 9 g Na $_{2}\text{HPO}_{4}$  x 12 H $_{2}$ 0

ReadyStream® media bags are gamma-irradiated (>25 kGray)





#### **Preparation**

Please strictly follow the recommendations in the ReadyStream<sup>®</sup> instrument user manual.

#### Storage

Store the media in the aluminum foil until use at +15 °C to +25 °C.

After reconstitution, store the ReadyStream<sup>®</sup> media bag up to 5 days at +15 °C to +25 °C, protect from UV light including sun light.

### **Quality Control**

- If the ReadyStream<sup>®</sup> media bag is removed from the instrument, place the ReadyStream<sup>®</sup> media bag with the tubing connector facing upwards. Do not stack filled ReadyStream<sup>®</sup> media bags on top of each other.
- After storage, mix to homogenize and visually check for the absence of contaminant.
- Filling and dispensing on 2 different instruments is not recommended. Never attempt to fill a ReadyStream<sup>®</sup> media bag twice.

Function	Control strains	Collection	Incubation	Method of control	Expected results
Productivity	Salmonella Typhimurium	ATCC 14028		Qualitative	Growth good to very good
	Salmonella Enteritidis	ATCC 13076	16 h - 20 h at 36 - 38 °C		
	Escherichia coli	ATCC 25922			
	Escherichia coli	ATCC 8739			
	Salmonella Abony	NCTC 6017	18 h - 24 h at 30 - 35 °C		
	Cronobacter sakazakii	ATCC 29544	16 h - 20 h at 34 - 38 °C		
	Cronobacter muytjensii	ATCC 51329	10 II - 20 II at 34 - 36 °C		
Diluent	Staphylococcus aureus	ATCC 25923		Quantitative	+/- 30% of original count
	Escherichia coli	ATCC 25922	45 - 60 min at room temperature		
	Escherichia coli	ATCC 8739			

Please refer to the actual batch related certificate of analysis.

The performance test is in accordance with the current version of EN ISO 11133.

#### **Ordering information**

Product	Packaging	Ordering No.
ReadyStream <sup>®</sup> Media Bag GranuCult <sup>®</sup> Buffered Peptone Water One bag for 30 L with a 3 L media bag (10x concentrated media)*	10x Bags with dehydrated media*	5.74846.0030
ReadyStream <sup>®</sup> Media Bag GranuCult <sup>®</sup> Buffered Peptone Water One bag for 100 L with a 10 L media bag (10x concentrated media)*	3x Bags with dehydrated media*	5.74846.0100
ReadyStream <sup>®</sup> Filter Set	5x Filter sets	5.74826.0001
ReadyStream <sup>®</sup> System	1x	RDYSTRM01

\*Bags are shipped with dry media who will be reconstituted with the ReadyStream<sup>®</sup> system. The 10x concentrated media allows to dispense up to 30 L of 1x media from the 3 L bag and up to 100 L of 1x media from the 10 L bag.

#### Note

The information in this document is subject to change without notice and should not be construed as a commitment by Merck KGaA, Darmstadt, Germany. Merck KGaA, Darmstadt, Germany assumes no responsibility for any errors that may appear in this document. This manual is believed to be complete and accurate at the time of publication. In no event shall Merck KGaA, Darmstadt, Germany be liable for incidental or consequential damage in connection with or arising from the use of this manual.

## To place an order or receive technical assistance

Order/Customer Service: SigmaAldrich.com/order

Technical Service: SigmaAldrich.com/techservice

SigmaAldrich.com/ReadyStream



Merck KGaA Frankfurter Strasse 250 64293 Darmstadt, Germany



© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. Merck, the vibrant M, Millipore, GranuCult and ReadyStream are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

Lit. No. MK\_DS8456EN Ver. 2.0 01/2022